

Ξ BARYONS

($S = -2, I = 1/2$)

$\Xi^0 = uss, \Xi^- = dss$

Ξ^0

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

Mass $m = 1314.83 \pm 0.20$ MeV

$m_{\Xi^-} - m_{\Xi^0} = 6.48 \pm 0.24$ MeV

Mean life $\tau = (2.90 \pm 0.09) \times 10^{-10}$ s

$c\tau = 8.71$ cm

Magnetic moment $\mu = -1.250 \pm 0.014 \mu_N$

Decay parameters

$\Lambda\pi^0 \quad \alpha = -0.411 \pm 0.022 \quad (S = 2.1)$

" $\phi = (21 \pm 12)^\circ$

" $\gamma = 0.85 [h]$

" $\Delta = (218_{-19}^{+12})^\circ [h]$


$\Lambda\gamma \quad \alpha = 0.4 \pm 0.4$

$\Sigma^0\gamma \quad \alpha = 0.20 \pm 0.32$

| Ξ^0 DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|--------------------------------|----------------------------------|-----------------------------------|----------------|
| $\Lambda\pi^0$ | $(99.51 \pm 0.05) \%$ | S=1.2 | 135 |
| $\Lambda\gamma$ | $(1.18 \pm 0.30) \times 10^{-3}$ | S=2.0 | 184 |
| $\Sigma^0\gamma$ | $(3.5 \pm 0.4) \times 10^{-3}$ | | 117 |
| $\Sigma^+ e^- \bar{\nu}_e$ | $(2.7 \pm 0.4) \times 10^{-4}$ | | 120 |
| $\Sigma^+ \mu^- \bar{\nu}_\mu$ | $< 1.1 \times 10^{-3}$ | CL=90% | 64 |

$\Delta S = \Delta Q$ (SQ) violating modes or $\Delta S = 2$ forbidden (S2) modes

| | | | | |
|--------------------------|----|------------------------|--------|-----|
| $\Sigma^- e^+ \nu_e$ | SQ | $< 9 \times 10^{-4}$ | CL=90% | 112 |
| $\Sigma^- \mu^+ \nu_\mu$ | SQ | $< 9 \times 10^{-4}$ | CL=90% | 49 |
| $p\pi^-$ | S2 | $< 4 \times 10^{-5}$ | CL=90% | 299 |
| $p e^- \bar{\nu}_e$ | S2 | $< 1.3 \times 10^{-3}$ | | 323 |
| $p \mu^- \bar{\nu}_\mu$ | S2 | $< 1.3 \times 10^{-3}$ | | 309 |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

$$\text{Mass } m = 1321.31 \pm 0.13 \text{ MeV}$$

$$\text{Mean life } \tau = (1.639 \pm 0.015) \times 10^{-10} \text{ s}$$

$$c\tau = 4.91 \text{ cm}$$

$$\text{Magnetic moment } \mu = -0.6507 \pm 0.0025 \mu_N$$

Decay parameters

$$\Lambda\pi^- \quad \alpha = -0.456 \pm 0.014 \quad (S = 1.8)$$

$$" \quad \phi = (4 \pm 4)^\circ$$

$$" \quad \gamma = 0.89 [h]$$

$$" \quad \Delta = (188 \pm 8)^\circ [h]$$

$$\Lambda e^- \bar{\nu}_e \quad g_A/g_V = -0.25 \pm 0.05 [f]$$

| Ξ^- DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | P (MeV/c) |
|--------------------------------|---|------------------|----------------|
| $\Lambda\pi^-$ | $(99.887 \pm 0.035) \%$ | | 139 |
| $\Sigma^- \gamma$ | $(1.27 \pm 0.23) \times 10^{-4}$ | | 118 |
| $\Lambda e^- \bar{\nu}_e$ | $(5.63 \pm 0.31) \times 10^{-4}$ | | 190 |
| $\Lambda\mu^- \bar{\nu}_\mu$ | $(3.5 \begin{smallmatrix} +3.5 \\ -2.2 \end{smallmatrix}) \times 10^{-4}$ | | 163 |
| $\Sigma^0 e^- \bar{\nu}_e$ | $(8.7 \pm 1.7) \times 10^{-5}$ | | 122 |
| $\Sigma^0 \mu^- \bar{\nu}_\mu$ | $< 8 \times 10^{-4}$ | 90% | 70 |
| $\Xi^0 e^- \bar{\nu}_e$ | $< 2.3 \times 10^{-3}$ | 90% | 6 |

$\Delta S = 2$ forbidden (S_2) modes

| | | | | | |
|------------------------------|-------|---------|------------------|-----|-----|
| $n\pi^-$ | S_2 | < 1.9 | $\times 10^{-5}$ | 90% | 303 |
| $ne^- \bar{\nu}_e$ | S_2 | < 3.2 | $\times 10^{-3}$ | 90% | 327 |
| $n\mu^- \bar{\nu}_\mu$ | S_2 | < 1.5 | % | 90% | 314 |
| $p\pi^- \pi^-$ | S_2 | < 4 | $\times 10^{-4}$ | 90% | 223 |
| $p\pi^- e^- \bar{\nu}_e$ | S_2 | < 4 | $\times 10^{-4}$ | 90% | 304 |
| $p\pi^- \mu^- \bar{\nu}_\mu$ | S_2 | < 4 | $\times 10^{-4}$ | 90% | 250 |
| $p\mu^- \mu^-$ | L | < 4 | $\times 10^{-4}$ | 90% | 272 |

$\Xi(1530) P_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$\Xi(1530)^0$ mass $m = 1531.80 \pm 0.32$ MeV ($S = 1.3$)

$\Xi(1530)^-$ mass $m = 1535.0 \pm 0.6$ MeV

$\Xi(1530)^0$ full width $\Gamma = 9.1 \pm 0.5$ MeV

$\Xi(1530)^-$ full width $\Gamma = 9.9^{+1.7}_{-1.9}$ MeV

| $\Xi(1530)$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|-------------------------|--------------------------------|------------------|-------------|
| $\Xi \pi$ | 100 % | | 152 |
| $\Xi \gamma$ | <4 % | 90% | 200 |

$\Xi(1690)$

$$I(J^P) = \frac{1}{2}(?^?)$$

Mass $m = 1690 \pm 10$ MeV [j]

Full width $\Gamma < 30$ MeV

| $\Xi(1690)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$ | seen | 240 |
| $\Sigma \bar{K}$ | seen | 51 |
| $\Xi \pi$ | seen | — |
| $\Xi^- \pi^+ \pi^-$ | possibly seen | 214 |

$\Xi(1820) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Mass $m = 1823 \pm 5$ MeV [j]

Full width $\Gamma = 24^{+15}_{-10}$ MeV [j]

| $\Xi(1820)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$ | large | 400 |
| $\Sigma \bar{K}$ | small | 320 |
| $\Xi \pi$ | small | 413 |
| $\Xi(1530) \pi$ | small | 234 |

$\Xi(1950)$

$$I(J^P) = \frac{1}{2}(??)$$

Mass $m = 1950 \pm 15$ MeV [j]
 Full width $\Gamma = 60 \pm 20$ MeV [j]

| $\Xi(1950)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $\Lambda \bar{K}$ | seen | 522 |
| $\Sigma \bar{K}$ | possibly seen | 460 |
| $\Xi \pi$ | seen | 518 |

$\Xi(2030)$

$$I(J^P) = \frac{1}{2}(\geq \frac{5}{2}?)$$

Mass $m = 2025 \pm 5$ MeV [j]
 Full width $\Gamma = 20^{+15}_{-5}$ MeV [j]

| $\Xi(2030)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $\Lambda \bar{K}$ | $\sim 20\%$ | 589 |
| $\Sigma \bar{K}$ | $\sim 80\%$ | 533 |
| $\Xi \pi$ | small | 573 |
| $\Xi(1530)\pi$ | small | 421 |
| $\Lambda \bar{K} \pi$ | small | 501 |
| $\Sigma \bar{K} \pi$ | small | 430 |